

The 114th iCeMS SEMINAR

CeMI Seminar Series 30

Fri 10 August 2012

10:30-12:00

How is a long strand of genomic DNA organized into a nucleus or chromosome?

Lecturer: **Prof. Kazuhiro Maeshima**

Biological Macromolecules Laboratory
National Institute of Genetics, Japan

Venue: Room #119 Research Building 1
Project Lab (#32) Kyoto University

How is 2-m-long genomic DNA organized into a mitotic chromosome or nucleus? The nucleosome fiber has long been assumed to be folded into a 30-nm chromatin fiber, and further helically folded larger fiber. However, when we observed frozen hydrated human mitotic cells using cryoelectron microscopy (cryo-EM), no higher-order structures including 30-nm chromatin fibers were found. To further investigate the bulk structure of mitotic chromosomes, we performed small-angle X-ray scattering (SAXS) at SPring-8. No structural feature larger than 11 nm was detected, even on a chromosome-diameter scale ($\sim 1 \mu\text{m}$). We also found a similar scattering pattern in interphase nuclei of HeLa cells in the range up to $\sim 275 \text{ nm}$. Our findings suggest a common structural feature in interphase and mitotic chromatins: compact and irregular folding of nucleosome fibers occurs without a 30-nm chromatin structure.

Contact: Harada Group at harada-g@icems.kyoto-u.ac.jp
Hosted by: iCeMS (Institute for Integrated Cell-Material Sciences), Kyoto University
Co-hosted by: Center for Frontier Medicine, Global COE Program, Kyoto University

